

# *The “pain matrix” in pain-free individuals*

Article

Accepted Version

Salomons, T. V., Iannetti, G. D., Liang, M. and Wood, J. N. (2016) The “pain matrix” in pain-free individuals. *JAMA Neurology*, 73 (6). pp. 755-756. ISSN 2168-6149 doi: <https://doi.org/10.1001/jamaneurol.2016.0653> Available at <https://centaur.reading.ac.uk/64895/>

It is advisable to refer to the publisher’s version if you intend to cite from the work. See [Guidance on citing](#).

Published version at: <http://dx.doi.org/10.1001/jamaneurol.2016.0653>

To link to this article DOI: <http://dx.doi.org/10.1001/jamaneurol.2016.0653>

Publisher: American Medical Association

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

[www.reading.ac.uk/centaur](http://www.reading.ac.uk/centaur)

**CentAUR**

Central Archive at the University of Reading

Reading’s research outputs online

# **The “Pain Matrix” in Pain-Free Individuals**

Tim V Salomons PhD

School of Psychology and Clinical Language Sciences, University of Reading, Reading RG6 7BE

UK [t.v.salomons@reading.ac.uk](mailto:t.v.salomons@reading.ac.uk)

Gian Domenico Iannetti MD, PhD

NPP, UCL, Gower Street, London WC1E 6BT UK n [g.iannetti@ucl.ac.uk](mailto:g.iannetti@ucl.ac.uk)

Meng Liang School of Medical Imaging, Tianjin Medical University, Tianjin, China.

1 Guangdong Road, Hexi District, Tianjin, 300203, China [mengliang.mic@gmail.com](mailto:mengliang.mic@gmail.com)

John N Wood PhD

Molecular Nociception Group, WIBR, UCL, Gower Street, London WC1E 6BT UK

[J.wood@ucl.ac.uk](mailto:J.wood@ucl.ac.uk)

All authors contributed equally to this study

591 words

**Corresponding Author**

[J.Wood@ucl.ac.uk](mailto:J.Wood@ucl.ac.uk)

**0044-2076796954**

## INTRODUCTION

Human functional imaging provides a correlative picture of brain activity during pain. A particular set of CNS structures (e.g. ACC, thalamus, insula) consistently respond to transient nociceptive stimuli causing pain. Activation of this so-called “pain matrix” or “pain signature” has been related to perceived pain intensity, both within and between-individuals<sup>1,2</sup> and is now considered a candidate biomarker for pain in medico-legal settings, as well as a tool for drug discovery. The pain-specific interpretation of such fMRI responses, although logically flawed<sup>3,4</sup>, remains pervasive. For example, a recent review states that “the most likely interpretation of activity in the pain matrix seems to be pain”<sup>5</sup>. Demonstrating the non-specificity of the “pain matrix” requires ruling out the presence of pain when highly-salient sensory stimuli are presented. Here we administer noxious mechanical stimuli to individuals with congenital insensitivity to pain and sample their brain activity with fMRI. Loss-of-function *SCN9A* mutations in these individuals fully impairs sodium channel Nav1.7 activity in peripheral neurons, resulting in loss of the ability to experience pain through impaired peripheral drive that leaves tactile percepts fully intact.<sup>5</sup> This allows complete experimental disambiguation of sensory responses and painful sensations.

## METHODS

3-Tesla fMRI was performed on two pain-free individuals (one female) and four age-matched controls. Subjects received twenty-four mechanical stimuli (465mN, 0.2mm tip, 1s duration) to their right hand dorsum. fMRI results from thermal stimuli are not reported due to motion artifacts. Subjects rated the intensity of both subjective sensation (0=no sensation, 10=most intense sensation imaginable) and pain (0=no pain, 10=most intense pain imaginable). GLM analysis of fMRI data were performed using FSL (<http://fsl.fmrib.ox.ac.uk/fsl>), using a cluster correction for multiple comparisons ( $z=1.96$ ,  $p<0.05$ ) at single-subject level and a conjunction analysis at group-level, such that group activations represent regions significantly activated in all individuals. To compare results to a canonical “pain matrix”, a meta analysis of pain studies ( $n=139$ ) was performed with Neurosynth ([www.Neurosynth.org](http://www.Neurosynth.org)) using forward inference with the feature set “painful”. Group comparisons were conducted by extracting activation z-scores from the Neurosynth-defined pain matrix and from key pain matrix regions (thalamus, insula, S2 and ACC - defined using the Harvard Oxford 25% probability atlas).

## RESULTS

In response to identical noxious stimuli, pain-free subjects reported similar levels of sensation to healthy controls [patients:  $4.6\pm0.5$ ; controls:  $4.4\pm1.2$  (mean $\pm$ SD),  $F=0.53$ ,  $p=0.51$ ]. Unlike controls, who uniformly reported the stimuli as painful ( $3.2\pm1.8$ ), the patients’ percepts were devoid of any painful quality. Strikingly, fMRI revealed normal activation of brain regions commonly activated by painful stimuli in both pain-free individuals (Figure 1a,c). There was no significant difference between patients and controls either across the entire “pain matrix” or in key “pain matrix” regions (Figure 1b; thalamus,

F=0.66, p=0.46; ACC, F=0.02, p=0.89; S@, F=0.01, p=0.93, Insula: F=0.09, p=0.78; pain matrix: F=0.3, p=0.61).

## DISCUSSION


Previous work<sup>3</sup> interpreting “pain matrix” activation as a response to salient sensory stimuli rather than perceptual qualities unique to pain has been challenged on the basis that the presence of pain in response to these stimuli could not be fully ruled out.<sup>5</sup> Here we address this challenge by demonstrating intact “pain matrix” responses in individuals congenitally unable to experience pain.

These observations reinforce the need for caution in using “pain matrix” responses for diagnosis or drug discovery and corroborate evidence that reported correlations between neuroimaging data and perceived pain have largely relied on non-pain-specific activities.<sup>43</sup> Examining how the brain gives rise to the unique perceptual experience of pain will require human neuroimaging to be supplemented by techniques that allow for causal inferences. These include studies in non-human species where cell populations and circuitry can be genetically or chemically modified, as well as human studies of individuals with relevant lesions or genetic mutations.


## Acknowledgements

The European Research Council (GDI), Wellcome Trust (JNW, GDI), MRC (JNW) and an EC Marie Curie Fellowship (TVS) supported this study, to which all authors contributed equally. We thank Professor Tom Johnstone for statistical consultation.


**A** Controls and neurosynth




**B** Cluster mean activation



**C** Pain-free patients and neurosynth



**D** Patients and controls conjunction



**FIGURE 1:** (A) shows the Neurosynth-based "pain matrix" (red) and the regions where all control subjects had significant activation in response to noxious stimulation (blue). (B) shows activation levels (z-scores) of single subjects within regions of the "pain matrix" (C) shows the Neurosynth-based "pain matrix" (red) and "pain matrix" regions where pain-free individuals had significant activation (yellow).

(D) shows the conjunction (green) of pain-free and control activations within the Neurosynth-based “pain matrix” regions.

## References

1. Wager TD, Atlas LY, Lindquist MA, Roy M, Woo CW, Kross E. An fMRI-based neurologic signature of physical pain. *N Engl J Med*. 2013 368(15):1388-97.
2. Huang G, Xiao P, Hung YS, Iannetti GD, Zhang ZG, Hu L. A novel approach to predict subjective pain perception from single-trial laser-evoked potentials. *Neuroimage*. 2013 ;81:283-93.
3. Iannetti GD, Salomons TV, Moayed M, Mouraux A, Davis KD. Beyond metaphor: contrasting mechanisms of social and physical pain. *Trends Cogn Sci*. 2013 17(8):371-8.
4. Hu L, Iannetti GD. Painful issues in pain prediction. *Trends Neurosci*. 2016, in press
5. Eisenberger NI. Social pain and the brain: controversies, questions, and where to go from here. *Annu Rev Psychol*. 2015 66:601-29.
6. Minett MS, Nassar MA, Clark AK, Passmore G, Dickenson AH, Wang F, Malcangio M, Wood JN. Distinct Nav1.7-dependent pain sensations require different sets of sensory and sympathetic neurons. *Nat Commun*. 2012 Apr 24;3:791